



Willow Lane
Community Primary School

Willow Lane CP School
Willow Lane,
Lancaster,
LA1 5PR
bursar@willow.lancs.sch.uk
0152465880

head@willow.lancs.sch.uk



Appletree Nursery School
Milking Stile Lane,
Lancaster,
LA1 5QB
bursar@appletree.lancs.sch.uk
0152464132

Head: Lucy Naylor

05/12/22

Dear Parent / Guardian

Re: Scarlet Fever

Please be aware that a small number of children are still being diagnosed with scarlet fever and Strep A.

We have sought advice from UK Health Security Agency North West Health Protection Team. They have advised that although scarlet fever is usually a mild illness, it should be treated with antibiotics to minimise the risk of complications and reduce the spread to others.

Symptoms of scarlet fever:

The **symptoms** of scarlet fever include a sore throat, headache, fever, nausea and vomiting. This is followed by a fine red rash which typically first appears on the chest and stomach, rapidly spreading to other parts of the body. On more darkly-pigmented skin, the scarlet rash may be harder to spot, but it should feel like 'sandpaper'. The face can be flushed red but pale around the mouth.

If you think you or your child have scarlet fever:

- See your GP or contact NHS 111 as soon as possible.
- Ask your GP to take a throat swab to confirm if this is scarlet fever.
- Make sure that you/your child takes the full course of any antibiotics prescribed by the doctor.
- Stay at home and away from nursery, school or work for **at least 24 hours after starting the antibiotic treatment**, to avoid spreading the infection.



If you/your child has an underlying condition which affects the immune system, you should contact your GP or hospital doctor to discuss whether any additional measures are needed.

Potential complications:

Children who have had **chickenpox** recently are more likely to develop more serious infection during an outbreak of scarlet fever and so parents/guardians should remain vigilant for symptoms such as a persistent high fever, cellulitis (skin infection) and arthritis (joint pain and swelling). If you are concerned for any reason please seek medical assistance immediately. **All children with chickenpox should remain off school / nursery until at least 5 days after the start of their rash AND until all their spots have crusted over.**

Infection control advice:

The Health Protection Team have advised us on hygiene measures to help prevent spread within our school / nursery. Parents/guardians can also help with this by reminding children of the importance of coughing / sneezing into tissues and placing these in the bin, washing hands regularly and ensuring any breaks to the skin are covered with a waterproof plaster/dressing before attending school / nursery.

Further information:

You can find more information at:

- NHS Choices <https://www.nhs.uk/conditions/scarlet-fever/>
- UKHSA Scarlet Fever Frequently Asked Questions
<https://www.gov.uk/government/publications/scarlet-fever-symptoms-diagnosis-treatment>

If you have any further questions, please contact the school office.

Yours sincerely,
Lucy Naylor

